HOW IMPORTANT IS IT TO YOU? | HOW READY ARE YOU? |

HOW CONFIDENT ARE YOU?

Tips for Giving Feedback







WORK

ALCOHOL









A Standard Drink

Any Drink Containing About 14 Grams Of Alcohol* *NIAAA (www.RethinkingDrinking.NIAAA.NIH.gov)

12 fl oz beer

5 fl oz table wine

1.5 fl oz liquor (vodka, tequila, etc.)







Craft beers often contain a higher % alcohol. See Lower Risk Drink Limits chart to know your limit.

Lower Risk Drink Limits*

	Per Day	Per Week
WOMEN	3	7
MEN	4	14
OVER 65	3	7
	LESS IS BETTER	

AVOID ALCOHOL IF YOU

- · take medications that interact with alcohol
- · have a health condition made worse by drinking
- · are under 21 years of age
- · plan to drive a vehicle or operate machinery
- · are pregnant or trying to become pregnant

O1 RAISE THE SUBJECT

Ask permission: "Would you mind taking a few minutes to discuss your screening results?"

02 **PROVIDE FEEDBACK**

- · Review reported alcohol use and refer to lower risk alcohol auidelines.
- Discuss possible health and other consequences of use; link to purpose of visit, if applicable.
- · Express concern.
- Elicit the person's response: "What do you think about this information?"

O3 ENHANCE MOTIVATION

- "On a scale of 0-10, how important is it to you to decrease (or quit) your drinking?"
- "On a scale of 0-10, how confident are you that you will be able to make this change?"
- · "How does your current level of drinking fit with what matters most to you?"
- · When readiness is low, ask, "What do you enjoy about drinking? What do you not enjoy about drinking?" Then summarize both sides.

04 NEGOTIATE AND ADVISE

- Elicit response: "What are your thoughts about our conversation?"
- · Negotiate a goal: "What steps are you interested in taking to make a change?"
- · Assist in developing a plan: "What could help you accomplish your goal? What will be challenging?"
- · Summarize the conversation. Arrange follow-up.
- Thank the person for having the conversation.

OPTIONS FOR MORE HELP Medication • Referral • www.LinkingCare.org This guide can be used for other risky behaviors, such as tobacco or illicit drug use. 8/2016





